



# Hybrid particle-field model for DNA

Sigbjørn Løland Bore  
Weekley Hylleraas seminar


10.05.2019

# Primer on DNA (1)



# Primer on DNA (2)

- ▶ Watson and Crick pairing
- ▶ Double helix formation
- ▶ Three types of helicies:
  - ▶ A-DNA
  - ▶ B-DNA
  - ▶ Z-DNA



# Primer on DNA (3)

- ▶ Human DNA: 1-3 m
- ▶ Persistence length: 50 nm
- ▶ Very flexible
- ▶ Environmental effects
  - ▶ Temperature
  - ▶ Salt




## The project:

- ▶ Reuse established CG-representations for DNA
- ▶ Model nonbonded interactions within hybrid particle-field framework
- ▶ Parametrize the model
- ▶ Benchmark the model
- ▶ No excuses, parallel implementation

# Coarse grain representation

- ▶ The coarse-grained representation should fulfill:
  - ▶ Represent the structural organization
  - ▶ 72 au per bead
- ▶ 3SPN-Model of Juan de Pablo
- ▶ Replace nonbonded interactions



# Bonded interactions

$$H_0(\{\mathbf{r}\}) = \sum_i^{N_{\text{atom}}} \frac{1}{2} m_i \dot{\mathbf{r}}_i^2 + \sum_i^{N_{\text{bond}}} \frac{1}{2} k_r (r_i - r_{i0})^2 + \sum_i^{N_{\text{bend}}} \frac{1}{2} k_\theta (\theta_i - \theta_{i0})^2 - \sum_i^{N_{\text{tor}}} k_\phi \exp \left[ -\frac{(\phi_i - \phi_{0i})^2}{2\sigma_\phi^2} \right],$$

Bond	$r_{i0}/\text{nm}$	Bend	$\theta_{i0}/\text{deg}$	Torsional	$\phi_{i0}/\text{deg}$
S-P	0.3899	S-P-S	94.49	P-S-P-S	-154.8
P-S	0.3559	P-S-P	120.15	S-P-S-P	-179.2
S-A	0.4670	A-S-P	112.07	A-S-P-S	-32.8
S-T	0.4189	P-S-A	103.53	S-P-S-A	54.8
S-G	0.4829	T-S-P	116.68	T-S-P-S	-44.8
S-C	0.3844	P-S-T	92.06	S-P-S-T	58.0
		G-S-P	110.12	G-S-P-S	-29.1
		P-S-G	107.40	S-P-S-G	53.9
		C-S-P	110.33	C-S-P-S	-34.1
		P-S-C	103.79	S-P-S-C	57.0

Equilibrium structure




# Hybrid particle field method

Mesoscale potentials in molecular dynamics:

$$V_{\text{ext},i} = \frac{1}{\tilde{\phi}_0} \left( k_b T \sum_j \chi_{ij} \phi_j(\mathbf{r}) + \frac{1}{\kappa} \left( \sum_j \phi_j(\mathbf{r}) - \tilde{\phi}_0 \right) \right)$$

$\chi_{ij}$ : Flory-Huggins parameter.  $\kappa$ : compressibility.  $\tilde{\phi}_0$ : system density.



$$\sum_{i < j} V_{ij}$$

$$\sum_i V(\{\phi(\mathbf{r}_i)\})$$



# Nonbonded interactions

$$W_{\text{elec}}[\rho] = \int d\mathbf{r} V_{\text{Coul}}(\mathbf{r})\rho(\mathbf{r})$$

$$W_{\text{non-elec}}[\{\phi\}] = \frac{1}{\tilde{\phi}_0} \int d\mathbf{r} \left[ \frac{k_b T}{2} \sum_{k,\ell} \chi_{k\ell} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r}) + \frac{1}{2\kappa} \left( \sum_k \phi_k(\mathbf{r}) - \tilde{\phi}_0 \right)^2 \right]$$

	P	S	A	T	C	G	W
P	$\chi_{PP}$	0	0	0	0	0	$\chi_{PW}$
S	0	0	0	0	0	0	0
A	0	0	0	$\chi_{NN}$	0	0	$\chi_{NW}$
T	0	0	$\chi_{NN}$	0	0	0	$\chi_{NW}$
C	0	0	0	0	0	$\chi_{NN}$	$\chi_{NW}$
G	0	0	0	0	$\chi_{NN}$	0	$\chi_{NW}$
W	$\chi_{PW}$	0	$\chi_{NW}$	$\chi_{NW}$	$\chi_{NW}$	$\chi_{NW}$	0


# Parametrization

- ▶ Parameters of the model:
  - ▶  $k_r, k_\theta, k_\phi, \chi_{NW}, \chi_{NN}, \chi_{PP}, \chi_{PW}$
- ▶ Goals:
  - ▶ Reproduces well the structure of B-DNA
  - ▶ Reproduce the persistence length of SS- and DS-DNA

# Optimization procedure(1): Fitness parameter

$$\eta = \frac{1}{N} \sqrt{\sum_{i=1}^N (\mathbf{r}_{i,1} - \mathbf{r}_{i,2})^2}$$


- ▶ Kabsch algorithm




## Optimization procedure(2): Optimization method

- ▶ Requirements:
  - ▶ No gradients
  - ▶ Handle noisy fitness
  - ▶ Few function calls

⇒ Bayesian Optimization






## Optimization procedure(3): Implementation



# Application of optimization

- ▶ 32 bp DNA, 100 mM salt
- ▶ 120ns




# Applications: Structural properties

► Best set:

- $\chi_{NW} = 19.0 \text{ kJ mol}^{-1}$
- $\chi_{NN} = -12.7 \text{ kJ mol}^{-1}$
- $\chi_{PW} = -7.2 \text{ kJ mol}^{-1}$
- $\chi_{PP} = -4.2 \text{ kJ mol}^{-1}$



Property	simulation	expt.
Bases per turn	$9.6 \pm 0.3$	10
Rise pr bp/nm	$0.34(5) \pm 0.01$	0.34
Radius/nm	$0.88 \pm 0.04$	0.94



# Applications: Hairpin-formation



Initial




Final




# Persistence length DS-DNA


$$\langle \hat{\mathbf{t}} \cdot \hat{\mathbf{t}}_l \rangle = e^{-l/l_p}, \quad \mathbf{t} \equiv \mathbf{r}_{P,i+10} - \mathbf{r}_{P,i}$$



- ▶ Experimental:  $l_p = 40\text{-}60$  nm
- ▶ Simulation:  $l_p = 43$  nm



# Problem with SS-DNA



Both are too stiff!

# Outlook

- ▶ Redo optimization to get better SS-strand behaviour
  - ▶ Less stiff  $k_\phi$
  - ▶ Limit  $\chi_{NW} \leq 10 \text{ kJ mol}^{-1}$
- ▶ Applications on longer doublestranded DNA
- ▶ Investigate the effect of salt on persistence length
- ▶ Plans for applying optimization on other systems

# Acknowledgements

Morten Ledum  
Michele Cascella



**OCCAM**  
Molecular Dynamics



**notur**



**Hylleraas**



**UiO** ●●